

TIGHT CLOSURE COMMUTES WITH LOCALIZATION IN BINOMIAL RINGS.

KAREN E. SMITH *

ABSTRACT. It is proved that tight closure commutes with localization in any domain which has a module finite extension in which tight closure is known to commute with localization. It follows that tight closure commutes with localization in binomial rings, in particular in semigroup or toric rings.

Tight closure, introduced in [HH1], is a closure operation performed on ideals in a commutative, Noetherian ring containing a field. Although it has many important applications to commutative algebra and related areas, some basic questions about tight closure remain open. One of the biggest such open problems is whether or not tight closure commutes with localization. For an exposition of tight closure, including the localization problem, see [Hu].

The purpose of this paper is give a simple proof that tight closure commutes with localization for a class of rings that includes semi-group rings and slightly more generally, the so-called "binomial" rings, that is, those that are a quotient of a finitely generated algebra by an ideal generated by binomials. Although this is a modest class of rings, the result here includes coordinate rings of all affine toric varieties, and is general enough to substantially improve previous results of a number a mathematicians, including Moty Katzman, Will Traves, Keith Pardue and Karen Chandler, Irena Swanson and myself. Thanks to Florian Enescu for pointing out a missing hypothesis in an earlier version.

Throughout this note, R denotes a Noetherian commutative ring of prime characteristic.

We say that *tight closure commutes with localization in R* if, for all ideals I in R and all multiplicative systems U in R , $I^*R[U^{-1}] = (IR[U^{-1}])^*$.

The main result is the following:

Theorem. *Let R be a ring with the following property: for each minimal prime P of R , the quotient R/P has a finite integral extension domain in which tight closure commutes with localization. Then tight closure commutes with localization in R .*

1991 *Mathematics Subject Classification.* 13A35.

Key words and phrases. tight closure, localization, binomial ideals, toric rings, semigroup rings.

* Supported by the National Science Foundation and the Alfred P. Sloan Foundation.

Lemma 1. *If tight closure commutes with localization in R/P for each minimal prime P of R , then tight closure commutes with localization in R .*

Proof. (See also [AHH 3.8]) It follows immediately from the definitions that $I^*R[U^{-1}] \subset (IR[U^{-1}])^*$ in general. To prove the reverse inclusion, let $\frac{z}{1}$ be in $(IR[U^{-1}])^*$. Then modulo each minimal prime of R disjoint from U , the image of $\frac{z}{1}$ is in $(IR/P[U^{-1}])^* = (IR/P)^*R/P[U^{-1}]$. For each minimal prime P_i disjoint from U , there is some u_i in U such that $u_i z$ is in $(IR/P_i)^*$. By setting $u = \prod u_i$, we see that $uz \in (IR/P)^*$ for all minimal primes P of R disjoint from U . On the other hand, for each minimal prime Q that meets U , pick any element in $Q \cap U$ and let v be their product. Now $uvz \in (IR/P)^*$ for every minimal prime P of R , which means that $uvz \in I^*$ in R . But because $uv \in U$, we conclude that $\frac{z}{1} \in I^*R[U^{-1}]$ and tight closure commutes with localization in R . $\square$

Lemma 2. *Let R be a domain that has a finite extension domain S in which tight closure commutes with localization. Then tight closure commutes with localization in R .*

Proof. Let $z \in R$ and let $I \subset R$. If $\frac{z}{1} \in (IR[U^{-1}])^*$, then the equations $c(\frac{z}{1})^q \in I^{[q]}R[U^{-1}]$ hold also after expansion to $S[U^{-1}]$, so that $\frac{z}{1}$ is in $(IS[U^{-1}])^*$. Since tight closure commutes with localization in S , we have $\frac{z}{1}$ is in $(IS)^*S[U^{-1}]$. This means that there exists a $u \in U$ such that $uz \in (IS)^*$. But in general it is easy to see that, $(IS)^* \cap R \subset I^*$ for any ideal I of R and any integral extension domain S of R , so $uz \in I^*$. (Indeed, by tensoring with the fraction field of R and then clearing denominators, one can find an R -linear map $S \xrightarrow{\phi} R$ which sends a fixed non-zero c in S to some non-zero element d of R . Then we can apply ϕ to the equations expressing $c(zu)^q \in I^{[q]}S$ to get equations expressing $d(uz)^q \in I^{[q]}$ in R . This implies that $\frac{z}{1} \in I^*R[U^{-1}]$ and the proof is complete. $\square$

Proof of Theorem. By the first lemma, it suffices to consider the case where R is a domain. The proof is complete by the second lemma. $\square$

Corollary. *If R has the property that modulo each minimal prime, the ring R/P has an integral extension that is F -regular, then tight closure commutes with localization in R .*

Proof. By definition, a ring is F -regular if all ideals are tightly closed both in the ring and in all its localizations. Thus tight closure trivially commutes with localization in an F -regular ring, and the corollary follows immediately from the Theorem. $\square$

Corollary. *Let R be a ring of the form $k[x_1, \dots, x_n]/J$ where J is generated by binomials in the x_i 's (a binomial is any polynomial of the form $\lambda x_1^{a_1} \dots x_n^{a_n} - \mu x_1^{b_1} \dots x_n^{b_n}$, where λ and μ are elements of k . In particular, every monomial is a binomial in this sense.) Then tight closure commutes with localization in R .*

Proof. There is no loss of generality in assuming the field is algebraically closed, by an argument similar to the one used to prove Lemma 2 (we can apply Lemma 2 directly if $\bar{k} \otimes_k R$ is a domain where $\bar{k}$ is the algebraic closure of k). In this case, the

minimal primes of J are themselves binomial ideals [ES], so by Lemma 1, we may assume that R is a domain. Now, a ring which is the quotient of a polynomial ring by a prime binomial ideal is isomorphic to a subring of a polynomial ring $k[t_1, t_2, \dots, t_d]$ generated by monomials in the t_i 's, so we assume R is such a monomial algebra [ES, 2.3]. The normalization $\tilde{R}$ of such a monomial algebra is again a monomial algebra. On the other hand, a normal monomial algebra is a direct summand of the polynomial overring $k[t_1, \dots, t_d]$ (see, *e.g.* [BH, Exercise 6.1.10]). Since a direct summand of a regular ring (or even F-regular ring) is F-regular, we see that $\tilde{R}$ is F-regular, so by the Corollary above, tight closure commutes with localization in R . $\square$

Special cases of this corollary, including the case where J is generated by monomials, had been observed using different methods by Will Traves, Moty Katzman, Keith Pardue and Karen Chandler, and Irena Swanson and myself. Katzman's work [K] includes an interesting approach to the localization problem using Gröbner basis in the special case of binomial ideals. Aldo Conca [C], and Karen Chandler and Keith Pardue, also made progress towards this Gröbner basis approach in the binomial ideal case. In [SS], a different approach to the localization problem is considered, in which the growth of the primary components of Frobenius powers is analyzed; the problem is solved for monomial ideals in monomial rings. In [T], the monomial case is solved using differential operators. The thesis of Doug McCulloch [Mc] does not directly address the localization problem, but contains a systemic study of tight closure in binomial rings.

REFERENCES

- [AHH1] Aberbach, I., Hochster, M. and Huneke, C., *Localization of tight closure and modules of finite phantom projective dimension*, J. Reine angew. **434** (1993), 67–114.
- [BH] Bruns, W. and Herzog, J., *Cohen-Macaulay Rings*, vol. 39, Cambridge Studies in advanced mathematics, 1993.
- [C] Conca, A., *Hilbert-Kunz function of monomial ideals and binomial hypersurfaces*, Manuscripta Math. **90** (1996), 287–300.
- [ES] Eisenbud, D. and Sturmfels, B., *Binomial ideals*, Duke **84** (1996), 1–45.
- [HH1] Hochster, M. and Huneke, C., *Tight closure, invariant theory, and the Briançon-Skoda theorem*, JAMS **3** (1990), 31–116.
- [Hu] Huneke, C., *Tight closure and its applications*, vol. 88, CBMS Conference Series, American Mathematical Society, Providence, RI, 1996.
- [K] Katzman, Mordechai, *The complexity of Frobenius powers of ideals.*, J. Algebra **203**, 211–225.
- [Mc] McCulloch, D., *Tight closure and base change*, thesis, University of Michigan (1997).
- [SS] Smith, K.E. and Swanson, I., *Linear growth of primary components of monomial ideals*, Communications in algebra **25** (1997), 3071–3079.
- [T] Traves, W., *Differential operators on monomial rings*, Jour. Pure and Applied Algebra **136** (1999), 183–197.

MATHEMATICS DEPARTMENT, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109